ELSEVIER

Environmental Pollution 150 (2007) I–III

ENVIRONMENTAL POLLUTION

ww.elsevier.com/locate/envpol

Contents of Volume 150

Number 1 2007

Editorial

1 **Environmental Pollution at volume 150**
W.J. Manning

2 **An analysis of papers published in Environmental Pollution in 2006: A continuing pattern of advancement and success**
Y.-G. Zhu, Z.-C. Wang, W.J. Manning

Reviews

5 **Occurrence, behavior and effects of nanoparticles in the environment**
B. Nowack, T.D. Bucheli

The behavior and the effects of natural and engineered nanoparticles in the environment are reviewed.

23 **Dynamics of trace metals in organisms and ecosystems: Prediction of metal bioconcentration in different organisms and estimation of exposure risks**
S. Fränzle, B. Markert, S. Wünschmann

Degradation, amounts and evolution of N-free vs. nitrogenous SOM control transfer of essential and toxic metals from soil into plants, to be estimated from coordination (bio-)chemistry.

34 **Phytoextraction of metals from soils: How far from practice?**
L. Van Nevel, J. Mertens, K. Oorts, K. Verheyen

Phytoextraction efficiency should be improved and associated risks need more attention before phytoextraction can be established as a commercial technology.

41 **Contaminant exposure in terrestrial vertebrates**
P.N. Smith, G.P. Cobb, C. Godard-Codding, D. Hoff, S.T. McMurry, T.R. Rainwater, K.D. Reynolds

Both biotic and abiotic factors determine chemical exposure for terrestrial vertebrates.

65 **Intercontinental transport of aerosols and photochemical oxidants from Asia and its consequences**
D.J. Wuebbles, H. Lei, J. Lin

Air quality over parts of North America is being affected by pollutants transported from Asia.

85 **Toward a biologically significant and usable standard for ozone that will also protect plants**
E. Paoletti, W.J. Manning

Current standards do not protect vegetation from ozone, but progress is being made.

96 **Increasing risk for negative ozone impacts on vegetation in northern Sweden**
P.E. Karlsson, L. Tang, J. Sundberg, D. Chen, A. Lindskog, H. Pleijel

Increasing risks for ozone impacts on vegetation in northern Sweden.

107 **Agricultural opportunities to mitigate greenhouse gas emissions**
J.M.-F. Johnson, A.J. Franzluebbers, S.L. Weyers, D.C. Reicosky

Management options can be used to reduce agriculture's environmental impacts.

125 **Challenges in quantifying biosphere–atmosphere exchange of nitrogen species**
M.A. Sutton, E. Nemitz, J.W. Erisman, C. Beier, K.B. Bahl, P. Cellier, W. de Vries, F. Cotrufo, U. Skiba, C. Di Marco, S. Jones, P. Laville, J.F. Soussana, B. Loubet, M. Twigg, D. Famulari, J. Whitehead, M.W. Gallagher, A. Neftel, C.R. Flechard, B. Herrmann, P.L. Calanca, J.K. Schjoerring, U. Daemmgen, L. Horvath, Y.S. Tang, B.A. Emmett, A. Tietema, J. Peñuelas, M. Kesik, N. Brueggemann, K. Pilegaard, T. Vesala, C.L. Campbell, J.E. Olesen, U. Dragosits, M.R. Theobald, P. Levy, D.C. Mobbs, R. Milne, N. Viovy, N. Vuichard, J.U. Smith, P. Smith, P. Bergamaschi, D. Fowler, S. Reis

Current N research is separated by form; the challenge is to link N components, scales and issues.

140 **Reduced nitrogen in ecology and the environment**
J.W. Erisman, A. Bleeker, J. Galloway, M.S. Sutton

Half of industrial ammonia production is eventually lost to the global environment with significant effects.

doi:10.1016/S0269-7491(07)00545-3

150 Global fate of POPs: Current and future research directions
R. Lohmann, K. Breivik, J. Dachs, D. Muir

Future studies into the global fate of POPs will need to pay more attention to the various biogeochemical and anthropogenic cycles to better understand emissions, transport and sinks.

166 Microbial interactions with organic contaminants in soil: Definitions, processes and measurement
K.T. Semple, K.J. Doick, L.Y. Wick, H. Harms

Understanding organic contaminant's behaviour in soil is key to chemically predicting biodegradation.

177 Accumulation of cyanobacterial toxins in freshwater "seafood" and its consequences for public health: A review
B.W. Ibelings, I. Chorus

Accumulation of cyanobacterial toxins in freshwater "seafood".

Number 2 2007

Regular Papers

193 Comparison of electrokinetic soil remediation methods using one fixed anode and approaching anodes
Z. Shen, X. Chen, J. Jia, L. Qu, W. Wang

During electrokinetic remediation, the nearer to the anode, the higher are the Cd removal velocities. A novel method with approaching anodes is proposed to accelerate remediation effect.

200 Use of ethylenediurea (EDU) to ameliorate ozone effects on purple coneflower (*Echinacea purpurea*)
Z. Szantoi, A.H. Chappelka, R.B. Muntifering, G.L. Somers

EDU can potentially ameliorate negative effects of O_3 on nutritive quality in purple coneflower.

209 Is an adjusted rhizosphere-based method valid for field assessment of metal phytoavailability? Application to non-contaminated soils
J. Fang, B. Wen, X.-q. Shan, J.-m. Lin, G. Owens

An adjusted-RHIZO method was the most promising approach for predicting metal phytoavailability to wheat under field conditions.

218 Short-term effects of dimethoate on metabolic responses in *Chrysolina pardalina* (Chrysomelidae) feeding on *Berkheya coddii* (Asteraceae), a hyper-accumulator of nickel
M. Augustyniak, P. Migula, J. Mesjasz-Przybyłowicz, M. Tarnawska, M. Nakonieczny, A. Babczyńska, W. Przybyłowicz, M.G. Augustyniak

Long-term exposure to high levels of nickel may have caused adaptive changes in the enzymes that enable Chrysolina pardalina to deal with other stressors, including organophosphate pesticides.

225 Poplar for the phytomanagement of boron contaminated sites
B.H. Robinson, S.R. Green, B. Chancerel, T.M. Mills, B.E. Clothier

Poplars reduce boron leaching from contaminated sites.

234 Temporal change in the distribution patterns of hexachlorobenzene and dichlorodiphenyltrichloroethane among various soil organic matter fractions
J.-j. Zhang, B. Wen, X.-q. Shan, S. Zhang, S.U. Khan

HCB and DDT tend to transfer from FA, HA and BHA fractions to lipid and IR fractions with increasing residence time.

243 Phytotoxicity of nanoparticles: Inhibition of seed germination and root growth
D. Lin, B. Xing

Engineered nanoparticles can inhibit the seed germination and root growth.

251 Adsorptive separation of cadmium from aqueous solutions and wastewaters by riverbed sand
Y.C. Sharma, S.N. Kaul, C.H. Weng

Riverbed sand for adsorptive separation of cadmium from waters and wastewaters.

258 Effects of ozone impact on the gas exchange and chlorophyll fluorescence of juvenile birch stems (*Betula pendula* Roth.)
C. Wittmann, R. Matyssek, H. Pfanz, M. Humar

Leaves of birch were more affected by ozone fumigation than corresponding stems, due to a higher ozone uptake rate.

267 Phytotoxic risk assessment of ambient air pollution on agricultural crops in Selangor State, Malaysia
S. Ishii, J.N.B. Bell, F.M. Marshall

There is a large potential of phytotoxic risk on vegetation in Selangor State, Malaysia.

280 Toxic effects of Pb^{2+} on growth of cowpea (*Vigna unguiculata*)
P.M. Kopittke, C.J. Asher, R.A. Kopittke, N.W. Menzies

The Pb^{2+} ion reduced the growth of cowpea by 10% at a solution activity of 0.2 μM for the shoots and 0.06 μM for the roots.

288 Variations of alkaline phosphatase activity and P fractions in sediments of a shallow Chinese eutrophic lake (Lake Taihu)
T. Zhang, X. Wang, X. Jin

Characteristics of the alkaline phosphatase activity and P fractions in sediments of different trophic status lake were studied in Lake Taihu.

Number 3 2007

Regular Papers

295 Physiological response of the lichen *Phaeophyscia hispidula* (Ach.) Essl., to the urban environment of Pauri and Srinagar (Garhwal), Himalayas, India
V. Shukla, D.K. Upreti

Lichen response due to ambient environmental changes.

300 Biological degradation of triclocarban and triclosan in a soil under aerobic and anaerobic conditions and comparison with environmental fate modelling
G.-G. Ying, X.-Y. Yu, R.S. Kookana

Triclocarban and triclosan can be degraded by microbial processes in aerobic soil, but will persist in anaerobic soil.

306 Acceleration of selenium volatilization in seleniferous agricultural drainage sediments amended with methionine and casein
G.S. Bañuelos, Z.-Q. Lin

Amending drainage sediment with either methionine or casein promotes the volatilization of selenium.

313 Modeling transport and fate of nitrogen from urea applied to a near-trench paddy field
X.Q. Liang, Y.X. Chen, H. Li, G.M. Tian, W.Z. Ni, M.M. He, Z.J. Zhang

N losses of near-trench paddy fields can be well predicted by water and N coupling model.

321 Nutrient flux in storm water runoff and baseflow from managed turf
K.W. King, J.C. Balogh, R.D. Harmel

Phosphorus losses from golf courses exceed levels to guard against eutrophication.

329 Year-to-year correlations in blood metal levels among individuals of two species of North American sea ducks
M. Wayland, R.T. Alisauskas, D. Kellett, J. Traylor, C. Swoboda, E. Neugebauer, K. Mehl

Some individuals of two species of sea ducks experience greater long-term (≥1 year) exposure to cadmium, selenium and lead compared to other individuals.

338 Effect of chemophytostabilization practices on arbuscular mycorrhiza colonization of *Deschampsia cespitosa* ecotype Waryński at different soil depths
E. Gucwa-Przepióra, E. Małkowski, A. Sas-Nowosielska, R. Kucharski, J. Krzyżak, A. Kita, P.F.A.M. Römkens

Addition of phosphorus and lignite in chemophytostabilization increased arbuscular mycorrhizal colonization of Deschampsia cespitosa roots.

347 Remediation of metal contaminated soil with mineral-amended composts
R. van Herwijnen, T.R. Hutchings, A. Al-Tabbaa, A.J. Moffat, M.L. Johns, S.K. Ouki

Composts can increase or decrease the bioavailability of metals in soil.

355 Leaf extracellular ascorbate in relation to O_3 tolerance of two soybean cultivars
F.-Y. Cheng, K.O. Burkey, J.M. Robinson, F.L. Booker

Leaf extracellular metabolites other than ascorbic acid are a factor in the differential ozone tolerance of two soybean cultivars.

363 Phytoextraction of cadmium and zinc from arable soils amended with sewage sludge using *Thlaspi caerulescens*: Development of a predictive model
A.P. Maxted, C.R. Black, H.M. West, N.M.J. Crout, S.P. McGrath, S.D. Young

A predictive model of Cd and Zn uptake by Thlaspi caerulescens is presented as a management tool in the phytoextraction of arable soils receiving sewage sludge.

373 Mercury and selenium in blue shark (*Prionace glauca*, L. 1758) and swordfish (*Xiphias gladius*, L. 1758) from two areas of the Atlantic Ocean
V. Branco, C. Vale, J. Canário, M.N. dos Santos

Mercury levels differ in Azores and Equator, and detoxification by selenium occurs.

Short Communication

381 Application and appreciation of chemical sand fixing agent-poly (aspartic acid) and its composites
J. Yang, H. Cao, F. Wang, T. Tan

The sand fixing agent was applied in powder form and it had no negative influence on plant growth.

I Contents of Volume 150